

SPLENECTOMY AND DYSFUNCTIONAL SPLEEN PROPHYLAXIS GUIDANCE IN ADULTS AND CHILDREN

Date effective from:	09/07/2025	Review date:	09/07/2028
Approved by:	Antimicrobial Management Committee (AMC) and NHS Lothian Clinical Management Group (CMG)		
Approval Date:	30/06/2025		
Author/s:	Antimicrobial Management Team (AMT) Pharmacists Carol Philip, Esperanza Palenzuela, and Sabrina Chan Antimicrobial Management Clinical Lead Simon Dewar		
Executive Lead:	AMT Lead Clinician		
Target Audience:	Medical, Surgical staff, anaesthetic staff, nursing teams and pharmacy team		
Supersedes:	Splenectomy and dysfunctional spleen prophylaxis guidance version 2.1		
Keywords (min. 5):	Surgical and medical prophylaxis, children adult, guideline, summary, antimicrobial, vaccine		

Version Control

Date	Author	Version/Page	Reason for Change
31/03/2022	Esperanza Palenzuela	Version 1.0	Immunisation updated, antibiotics dose frequency terminology and updated some links
09/01/2023	Esperanza Palenzuela	2.0 Page 4	Updated link for Immunisation and coeliac disease
11/01/2023	Esperanza Palenzuela	2.0 Page 8	Spelling corrected for MenACWY
07/02/2023	Esperanza Palenzuela	2.1 Page 4	Clarification of antibiotic prophylaxis in coeliac patients with dysfunctional spleen
08/07/2025	Sabrina Chan	3.0	Added advice on COVID-19 vaccination; updated criteria for lifelong antibiotic prophylaxis as per updated BSH guidance in 2024; updated link for travel advice; added link to antibiotic guidance for animal bites; updated discharge checklist.
24/09/2025	Sabrina Chan	3.1	Clarify wording regarding asplenia/ hyposplenism. Ensure consistency throughout document
28/10/2025	Esperanza Palenzuela	3.1	Add "MenQuadfi" new brand of MenACWY quadrivalent conjugate vaccine.

CONTENTS

Introduction	4
Risks of Absent or Dysfunctional Spleen	4
Vaccination	4
Elective Splenectomy	4
Emergency Splenectomy.....	4
Child Antibiotic Prophylaxis	8
Adult Antibiotic Prophylaxis	9
Rescue antibiotic therapy	10
Children rescue treatment – provide 5 days supply	10
Adult rescue treatment provide 5 days supply	10
Special patients groups and additional information	11
Chemotherapy, Radiotherapy or other Immunosuppressive treatment.....	11
Pregnancy/Breast-Feeding.....	12
1. Vaccines	12
2. ANTIBIOTICS	12
Travel	12
Animal bites.....	12
Tick bites	12
Clinical management and discharge checklist	13
References	14

INTRODUCTION

This guideline gives recommendations for the prevention of infection in patients who have an absent spleen/undergone a splenectomy (asplenia) or who have been diagnosed with a dysfunctional spleen (hyposplenism).

Splenectomy (Asplenia): Elective: e.g., for haematological disease or splenic abscess, cysts, mass, and neoplasm. **Emergency:** e.g., for traumatic injury to spleen or intra operative splenic injury.

Dysfunctional Spleen (Hyposplenism): e.g., conditions such as homozygous sickle cell disease and coeliac disease may lead to splenic dysfunction.

Immunisation is recommended in all patients with coeliac disease. The Lothian consensus guidance [Immunisation and coeliac disease](#) from NHS Lothian Coeliac Services and Lothian Immunisation Co-ordinating Group should be followed for vaccines.

Coeliac patients with a splenectomy require antibiotic prophylaxis. In coeliac patients with functional hyposplenism there is less consensus on the need for antibiotic prophylaxis and it is not routinely given.

RISKS OF ABSENT OR DYSFUNCTIONAL SPLEEN

Overwhelming infection is a major risk in patients with an absent (asplenia) or dysfunctional spleen (hyposplenism) and although uncommon, is associated with a high mortality. These infections are often due to encapsulated bacteria such as *Streptococcus pneumonia*, *Haemophilus influenzae* type B and *Neisseria meningitides*, more than half of those infected die. Other serious infections include malaria, babesiosis (caused by tick bite) and *Capnocytophaga canimorsus* (caused by dog bites) and secondary infections following influenza. For these reasons, it is imperative that all patients with an absent or dysfunctional spleen are appropriately immunized, receive appropriate antibiotics and counselling.

VACCINATION

ELECTIVE SPLENECTOMY

Start immunisation course at least TWO weeks **prior** to surgery (ideally four to six weeks).

EMERGENCY SPLENECTOMY

Start immunisation course at least TWO weeks **post**-surgery. However, if the patient is discharged earlier than this, they should start their immunisation schedule immediately before discharge. However, functional antibody response is better with delayed 14 days vaccination.

Given the changing pattern of routine vaccination, patients of different ages may have different “routine” vaccination histories. It is therefore essential to assess vaccination requirements against an individual’s vaccination history.

Check patient has been vaccinated according to UK schedule:

www.gov.uk/government/publications/the-complete-routine-immunisation-schedule

VACCINES FOR FIRST DIAGNOSED UNDER 1 YEAR OF AGE

Children should be fully immunized according to the national schedule, and should also receive

- Two doses of **MenACWY** vaccine at least one month apart during their first year
- An additional priming dose of **PCV13**, such as to receive a total of two priming doses at least 8 weeks apart commencing no earlier than 6 weeks of age in their first year. One additional booster dose of **PCV13** and one dose of **MenACWY** conjugate vaccine 8 weeks after the vaccinations scheduled at 1 year of age
- One dose of **PPV23** after the second birthday and at least 8 weeks after the last dose of **PCV13**
- Annual **influenza** vaccine each season for patients aged over 6 months

FIRST DIAGNOSIS AT 12-23 MONTHS OF AGE

If not yet administered, give the routine 1 year of age vaccines: **Hib/MenC**, **PCV13**, **MMR** and **MenB**, plus

- One additional dose of **PCV13** and one dose of **MenACWY** conjugate vaccine 8 weeks after the vaccinations scheduled at 1 year of age and
- One dose of **PPV23** after the second birthday and at least 8 weeks after the last dose of **PCV13**
- Annual **influenza** vaccine each season

FIRST DIAGNOSED FROM TWO YEARS TO UNDER TEN YEARS OF AGE

Ensure children are immunized according to the national schedule, and they should also receive:

- One dose of **PPV23**, followed by
- One dose of **MenACWY** conjugate vaccine
- If not already received the routine 2+1 schedule for **MenB**, ensure they have received two doses of **MenB** 8 weeks apart since first birthday
- If they have not received any **PCV** previously, they should receive a dose of this first followed by the dose of **PPV23** at least 8 weeks later
- Annual **influenza** vaccine each season

FIRST DIAGNOSED AT AGE TEN YEARS ONWARDS
<p>Older children and adults, regardless of previous vaccination, should receive:</p> <ul style="list-style-type: none"> • One dose of PPV23, MenB and MenACWY conjugate followed by • One additional MenB vaccine dose 4 weeks later • Annual influenza vaccine each season
VACCINATION SCHEDULE
<ul style="list-style-type: none"> • Offer annual influenza vaccine to all patients • Boosters of PPV23 vaccination every 5 years
VACCINES TERMINOLOGY
<p>MenACWY Conjugate = Meningococcal A, C, W135 and Y conjugate vaccine (Menveo, Nimenrix, MenQuadfi)</p> <p>Men B = Meningococcal B vaccine (Bexsero)</p> <p>Hib/Men C = Haemophilus type B conjugate vaccine (Menitorix)</p> <p>PCV13 = Pneumococcal conjugate vaccine (Prevenar 13)</p> <p>PPV23 = Pneumococcal polysaccharide vaccine</p> <p>MMR = Measles, mumps, and Rubella vaccine (Priorix)</p>
CORONAVIRUS (COVID-19) VACCINATION
<p>See the latest national guidance on COVID-19 vaccination programme, Green Book Chapter 14a COVID-19: the green book, chapter 14a - GOV.UK and Public Health Scotland Scottish Vaccine Update https://publichealthscotland.scot/publications/scottish-vaccine-update.</p>

ANTIBIOTIC PROPHYLAXIS

The increased risk of infections in patients with asplenia/hyposplenism is lifelong, but they are at the highest risk of infection and overwhelming sepsis in the immediate post-splenectomy period and for 1 to 3 years after. Therefore, antibiotic prophylaxis should be started immediately post-operatively and continue for at least 2 years.

Children less than 5 years should continue antibiotic prophylaxis until at least 5 years of age, and for a minimum of 2 complete years following splenectomy. For example, if a child had splenectomy at the age of 2, they should continue antibiotic prophylaxis until at least 5 years old. If they had splenectomy at the age of 4, they should continue antibiotic prophylaxis until at least 6 years old.

Penicillin prophylaxis is highly effective in children with sickle cell disease.

Long-term prophylactic antibiotics, beyond 2 years post-splenectomy, should be offered to patients considered at continued high risk of pneumococcal infection.

Factors associated with high risk of pneumococcal infections in asplenia/hyposplenism include:

- Extremes of age (< 5 years # or > 65 years old*).
- A history of previous invasive pneumococcal disease (IPD).
- Treated for underlying haematological malignancy, particularly those who have received splenic irradiation or ongoing immunosuppression.
- Patients with active ongoing graft-versus-host disease (GvHD).

Patient with the above risk factors will also need careful counselling and follow-up to ensure adherence to antibiotic prophylaxis.

The use of life-long antibiotics has potential disadvantages, such as the development of bacterial resistance, potential side effects, and poor adherence. Therefore, after the first 2 years post-splenectomy, the prescriber should review if the patient remains at high risk and if long-term prophylaxis is still beneficial.

Patients no longer at high risk after the first 2 years should be counselled regarding the risks and benefits of long-term antibiotics, and may choose to continue or discontinue prophylaxis. If chosen to stop long-term antibiotic prophylaxis, appropriate rescue antibiotic therapy should still be given to the patient for emergency use (see next section on Rescue Antibiotic Therapy).

*For asplenic/hyposplenic patients above 65 years of age and had previously discontinued their long-term antibiotic prophylaxis, prescribers should offer to restart long-term antibiotic prophylaxis.

CHILD ANTIBIOTIC PROPHYLAXIS		
	Prophylaxis	Duration
First line	<p>Phenoxymethylpenicillin (Penicillin V):</p> <p>Under 1 year; 62.5mg every 12 hours, orally</p> <p>1 – 4 years; 125mg every 12 hours, orally</p> <p>5 – 17 years; 250mg every 12 hours, orally</p> <p>If cover also needed for <i>H.influenzae</i> (e.g. incomplete vaccination) in child give Amoxicillin instead:</p> <p>1 month – 4 years; 125mg every 12 hours, orally</p> <p>5 – 11 years; 250mg every 12 hours, orally</p> <p>12 – 17 years; 500mg every 12 hours, orally</p>	Antibiotic prophylaxis should be continued until at least 5 years of age, and for a minimum of 2 years.
If penicillin allergy	<p>Erythromycin:</p> <p>1 month – 2 years; 125mg every 12 hours, orally</p> <p>2 – 7 years; 250mg every 12 hours, orally</p> <p>8 – 17 years; 500mg every 12 hours, orally</p>	
If nil-by-mouth following splenectomy	<p>Benzylpenicillin:</p> <p>1 month – 17 years; 25mg/kg every 12 hours, intravenously</p>	Additional cover with IV Benzylpenicillin is not required if the patient is already receiving antibiotics with appropriate activity (e.g., amoxicillin, cephalosporins or other beta-lactams). If unsure or patient is allergic to penicillin discuss with Microbiology.

ADULT ANTIBIOTIC PROPHYLAXIS		
	Prophylaxis	Duration
First line	Phenoxymethylpenicillin (Penicillin V): 250mg every 12 hours, orally	Minimum 2 years; lifelong preferred if considered high risk (see criteria on page 7)
If penicillin allergy	Clarithromycin 250mg every 12 hours, orally	
If nil-by-mouth following splenectomy	Benzylpenicillin 1.2g every 12 hours, intravenously	Additional cover with IV Benzylpenicillin is not required if the patient is already receiving antibiotics with appropriate activity (e.g., amoxicillin, cephalosporins or other beta-lactams). If unsure or patient is allergic to penicillin discuss with Microbiology.

RESCUE ANTIBIOTIC THERAPY

Patients may develop infection despite vaccination and antimicrobial prophylaxis. An emergency supply of oral antibiotics should be given to **all** the patients to keep at home - advise them to replace before the expiry dates.

Rescue antibiotics can be used however it is vital that patients seek immediate medical attention if they are unwell with symptoms in-keeping with infection e.g., raised temperature, malaise, or shivering.

For patients already taking antibiotic prophylaxis, the rescue antibiotic should be from a different antibiotic class to minimise possibilities of bacterial resistance. Choice of antibiotic should be made with regard to appropriate microbiology advice and local guidelines.

CHILDREN RESCUE TREATMENT – PROVIDE 5 DAYS SUPPLY	
First line:	Amoxicillin: Child 1 month -11 months ; 125mg every 8 hours, orally Child 1 - 4 years ; 250mg every 8 hours, orally Child 5 - 11 years ; 500mg every 8 hours, orally Child 12 - 17 years ; 500mg every 8 hours, orally
If penicillin allergy	Erythromycin: Child 2 - 7 years ; 250mg every 6 hours, orally Child 8 - 17 years ; 500mg every 6 hours, orally

ADULT RESCUE TREATMENT PROVIDE 5 DAYS SUPPLY	
First line:	Amoxicillin 500mg every 8 hours, orally
If penicillin allergy	Clarithromycin 500mg every 12 hours, orally

CHEMOTHERAPY, RADIOTHERAPY OR OTHER IMMUNOSUPPRESSIVE TREATMENT

- Most vaccines used post-splenectomy (see “Vaccination” section above) are inactivated – MenACWY, PCV13, PPV23, Hib/MenC, MenB, Influenza – with the exception of Fluenz Tetra and MMR vaccine which are live vaccines. The inactivated vaccines cannot replicate and can be administered to immunosuppressed individuals, although they may be a lower response than in an immuno-competent individual.
- Vaccinations should be given at least two weeks (ideally 4 -6 weeks) before initiation of treatment such as chemotherapy or radiotherapy. Where it is not possible vaccination beforehand, splenectomy, chemotherapy or radiotherapy should never be delayed.
- If not practicable to vaccinate two weeks before the initiation of chemotherapy and/or radiotherapy, immunisation can be delayed until at least three months after completion of therapy to maximise the response to the vaccine, whilst ensuring adequate antibiotic cover is prescribed in the interim.
- Individuals with immunosuppression should be vaccinated in accordance with the standard schedule but it should be borne in mind that these individuals may not make full antibody response.
- The NHS Lothian Oncology Online Quality System (OOQS) provides guidance and information for patients and carers following splenectomy (from NHS inform) [Patient information on vaccination post splenectomy](#)
- Patients who have undergone allogeneic bone marrow transplant* have functional hyposplenism. These patients require specific re-vaccination according to a separate policy, as per the National Allogeneic Bone Marrow Transplant Centre at the QEH, Glasgow.

This is available at: [Scottish Haematology Society | SHS Guidelines](#)

*(i.e., using stem cells from a bone marrow donor, including related and unrelated donors)

PREGNANCY/BREAST-FEEDING

1. VACCINES

Influenza vaccine should be given to all before the flu season, regardless of the stage of pregnancy.

Pneumococcal, Haemophilus and Meningococcal vaccine should be given when protection is required without delay.

Pertussis vaccine (now comes in the form of ADACEL® - tetanus, diphtheria, pertussis vaccine (Tdap)) should be offered to all women from 16 weeks of pregnancy, ideally by 32 weeks but beneficial until 38 weeks gestation.

There is no evidence of risk from vaccinating pregnant women or those who are breast feeding with inactivated viral or bacterial vaccine or toxoids.

2. ANTIBIOTICS

Penicillin's may be used in pregnancy if clinically indicated. However, other risk factors may be present in individual cases which may independently increase the risk of adverse pregnancy outcome. Clinicians are reminded to perform a case-specific risk assessment.

Do not suggest Erythromycin as there is an increased overall risk of congenital malformation or cardiac malformation.

Further information available from:

- <https://www.toxbase.org>
- <https://www.sps.nhs.uk/articles/safety-in-lactation-macrolides/>

TRAVEL

Patients with an absent or dysfunctional spleen are at risk of severe falciparum malaria. Guidance should be given on appropriate malaria prophylaxis and the need for close adherence to it. They are also at an increased risk of meningitis and other travel associated infections. Please check for destination-specific risks and disease prevention information, including any vaccine recommendations, on the Travel Health Pro Website [NaTHNaC - Asplenia and hyposplenia](#).

ANIMAL BITES

All animal bites need to be treated quickly, to reduce the chance of infection from *Capnocytophaga canimorsus*, which can lead to fulminant sepsis. All animal bites require immediate medical attention and may need antibiotic treatment – see NHS Lothian antimicrobial prescribing guideline on [Human and animal bites | Right Decisions](#).

INSECTS AND TICK BITES

Babesiosis is a rare tick-borne infection that can cause moderate to severe disease, including haemolytic anaemias. Therefore, it is essential to take precautions against being bitten in endemic areas.

CLINICAL MANAGEMENT AND DISCHARGE CHECKLIST

Appropriate vaccinations given	<input type="checkbox"/>
Appropriate antibiotic prophylaxis prescribed	<input type="checkbox"/>
Given course of emergency antibiotics on discharge?	<input type="checkbox"/>
<p>Patients should receive a "no functioning spleen" alert card and leaflet* "A guide for people without a working spleen", available from Health Protection Team: https://publichealthscotland.scot/publications/a-guide-for-people-without-a-working-spleen/</p> <p><i>*Note the PHS leaflet has not been updated on the Antibiotics page – i.e. the recommended age for long-term antibiotic prophylaxis is now < 5 years instead of < 16 years.</i></p> <p>Health Directorates: ☎ 0131 244 2241 or email: immunisationprogrammes@gov.scot</p>	<input type="checkbox"/>
Advise patients that they may wish to invest in an alert bracelet/tag	<input type="checkbox"/>
Discuss annual influenza vaccine and Pneumococcal vaccine every 5 years	<input type="checkbox"/>
Discuss need for immediate medical attention following animal bite and further vaccines if travelling and malaria precautions	<input type="checkbox"/>
Immediate discharge letter (IDL) informs GP that a splenectomy has been performed	<input type="checkbox"/>
IDL includes advice for GP to review need for long-term antibiotic prophylaxis after 2 years, or if life-long antibiotic prophylaxis is indicated.	<input type="checkbox"/>
Signpost patients to NHS inform https://www.nhsinform.scot/healthy-living/immunisation/when-to-immunise/children-and-adults-without-a-spleen-asplenia	<input type="checkbox"/>

REFERENCES

1. Prevention and treatment of infection in patients with an absent or hypofunctional spleen: A British Society for Haematology Guideline. *Br J Haematol.* 2024; 204:1672-1686. [Online] [British Journal of Haematology | Wiley Online Library](#) (Accessed May 2025)
2. Green Book chapter 7 at <https://www.gov.uk/government/publications/immunisation-of-individuals-with-underlying-medical-conditions-the-green-book-chapter-7> Last updated January 2020 (Accessed May 2025)
3. Green Book chapter 14a at <https://www.gov.uk/government/publications/covid-19-the-green-book-chapter-14a>. Last updated March 2022.(Accessed May 2025)
4. Green Book chapter 6 at https://assets.publishing.service.gov.uk/media/5a82ce28e5274a2e8ab5970f/Greenbook_chapter_6.pdf Last updated August 2017 (Accessed May 2025)
5. GOV.UK: Pertussis (whooping cough) vaccination programme for pregnant women: information for healthcare practitioners <https://www.gov.uk/government/publications/vaccination-against-pertussis-whooping-cough-for-pregnant-women/pertussis-whooping-cough-vaccination-programme-for-pregnant-women> Last updated August 2024 (Accessed June 2025)
6. Trust Guidelines for Patients with an Absent or Dysfunctional Spleen- Norfolk and Norwich University Hospitals (Accessed April 2023)
7. Spleen problems and spleen removal. <https://www.nhs.uk/conditions/spleen-problems-and-spleen-removal/> (Accessed February 2022)
8. Children and adults without spleen (asplenia). <https://www.nhsinform.scot/healthy-living/immunisation/when-to-immunise/children-and-adults-without-a-spleen-asplenia> (Accessed February 2022)
9. Preventing and treating infection in Children with asplenia or hyposplenia. *Paediatric Child Health* 2014;19(5):271-274 (Accessed February 2022)
10. Lothian Medicines Information Service Enquiry No 6688. Date completed 22/10/2014
11. Antibiotic prophylaxis and dysfunctional hyposplenism in coeliac disease. <https://gut.bmj.com/content/gutjnl/63/8/1210.full.pdf> (Accessed February 2023)
12. Antibiotic prophylaxis and dysfunctional hyposplenism in coeliac disease <https://cks.nice.org.uk/topics/coeliac-disease/management/management-of-confirmed-coeliac-disease/> (Accessed February 2023)
13. Antibiotic prophylaxis and dysfunctional hyposplenism in coeliac disease <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4103047/>(Accessed February 2023)